

Research Article

Phytoplankton assemblages in the ballast water of the navigated ships from the Indian Ocean to Shahid Rajaei Port, Hormozgan, Persian Gulf, Iran

Amidi R.¹; Fatemi S.M.R.^{2*}; Ghodousi J.³; Javid A.H.⁴

Received: April 2021

Accepted: August 2021

Abstract

The present study has been conducted to identify and enumerate ballast water phytoplankton on 24 ships entering Shahid Rajaei Port (13 bulk carriers, 4 oil tankers, 6 general cargo and 1 container), in two cold and warm seasons in winter 2017 and summer 2018, from the four origins as the name the Persian Gulf, western, eastern, and northwestern Indian Ocean. The sampling of phytoplankton was conducted by passing 100 liters of sampling waters through a 20-micron plankton net, in compliance with the G2 standard method approved by MEPC.173 (58). In this manner, 108 phytoplankton species were identified in the ballast water tanks of the sampled ships, belonging to 4 phylum, 5 classes, 23 orders, and 38 families. In the warm season, Dinophyceae (54%) and in the cold season, Bacillariophyceae (60.96%) were prevalent among the samples. The identified species from the Persian Gulf had the highest similarity with the species from the western Indian Ocean, and the least similarity with the eastern Indian Ocean. Apart from *Odontella longicruris*, all other species had been reported in the waters of the Persian Gulf in previous studies. This result shows that the phytoplankton assemblages between Shahid Rajaei Port and other parts of the Indian Ocean have a high similarity and does not impose a real threat to the environment from the ballast water transfer. However, from the identified species, 28 species (25.9%) are known to cause red tide and death of other aquatic species in the region, which highlights the importance of continuous monitoring and discharge control in the marine environment of the region.

Keywords: Phytoplankton, Invasive species, Ballast water, Shahid Rajaei port, Persian Gulf, Iran

1-Department of Environmental Science, Faculty of Natural Resources and Environment, Science and Research Branch, Islamic Azad University, Tehran, Iran

2-Department of Marine Science, Faculty of Natural Resources and Environment, Science and Research Branch, Islamic Azad University, Tehran, Iran

3-Department of Environmental Management, Faculty of Natural Resources and Environment, Science and Research Branch, Islamic Azad University, Tehran, Iran

4-Department of Environmental Engineering, Faculty of Natural Resources and Environment, Science and Research Branch, Islamic Azad University, Tehran, Iran

*Corresponding author's Email: fatemi_re@srbiau.ac.ir

Introduction

One of the four major threats against oceans, and the main cause of transfer for 40 percent of non-indigenous and invasive species to the ports around the world, is the unintentional transfer of these species through ballast water (Hallegraeff and Bolch, 1992; Ruiz *et al.*, 1997; Dickman and Zhang, 1999; Endresen *et al.*, 2004; David *et al.*, 2007; Pereira *et al.*, 2014; Steichen *et al.*, 2014; Soler-Figueroa *et al.*, 2020). The growth of commerce, expansion of shipping routes, increase in speed, traffic, capacity, and the number of ships on the one hand, and reduced length of voyages, lower number of maritime incidents, promoted safety of navigation, and carriage of 3 to 5 billion tons of ballast water around the world per year (Drake *et al.*, 2007; Tsolaki and Diamadopoulos, 2010; Satir, 2014; Cordell *et al.*, 2015) on the otherhand has resulted in the transfer of 7 to 10 thousand living species from one location to another (Drake and Lodge, 2004; Lakshmi *et al.*, 2021), in a manner that every 9 weeks, an invasive species is relocated to a new environment (Dickman and Zhang, 1999; Steichen *et al.*, 2014). The entry of invasive species into an ecosystem, with the assumption of their stability and persistence in the new environment and threat toward the indigenous species, imposes two major issues on the region: 1) impact on the economic and fisheries species, which would, in turn, affect the revenues generated from fishing activities; and 2) jeopardized biodiversity of the region, through

threatening one or more indigenous species, which will leave a long-term impact, as elimination of a species from the life cycle will affect the related cycles before and after that species, and the whole ecosystem as a result (Burkholder *et al.*, 2007; Balaji *et al.*, 2014; de Castro *et al.*, 2017).

To control the transfer and invasion by non-indigenous aquatic species through the ballast water of ships, the International Maritime Organization (IMO) responsible for developing international regulations for the safety of navigation and marine environment protection, has defined an approved ballast water management methods in the form of the International Convention for Control and Management of Ships' Ballast Water and Sediments, for all ships above 400 GT (Gross tonnage) following their age and ballast water capacity (Gollasch *et al.*, 2007; Tjahjono *et al.*, 2017; Čampara *et al.*, 2019). This convention entered into force on 8 September 2017, the Islamic Republic of Iran ratified the convention as the 28th party (Balley and Munawar, 2015).

According to Regulation B-4 of the convention, all ships shall exchange their ballast water at least 200 nautical miles from the nearest land and in the water at least 200 meters in depth. If not possible for any reason, the exchange should be as far from the nearest land as possible, and in all cases at least 50 nautical miles from the nearest land and in the water at least 200 meters in depth, or within oceanic waters, due to the different conditions between coastal

and oceanic areas (Cordell *et al.*, 2015). As per Resolution MEPC.297 (72) regarding revision of Regulation B-3, all tankers from 150 to 400 GT, and other ships not required to have IOPP Certificates shall install ballast water management systems onboard complying with Regulation D-2 not later than 8 September 2024 (Gerhard *et al.*, 2019; Van Slooten, 2021). By the same token, the Marine Environment Protection Committee (MEPC) identified the Persian Gulf as a special area in 2007 (MEPC.1/Circ.778/Rev.3), which highlights the particular oceanic conditions, extensive traffic of ships, and the environmental sensitivity and vulnerability of the region to exert special measures over maritime activities, especially shipping (Wang and Liu, 2021).

The Persian Gulf is a semi-enclosed, secluded, and shallow sea (Al-Majed, 2000), and is considered among the most diverse marine ecosystems with its low depth (average 35m), high salinity (about 40ppt), and high temperature (average 26°C), and low precipitation (below 100mm). It houses rich reserves of aquatic species and marine biodiversity of fish, and the majority is of native variety and low tolerance range against environmental changes due to severe stresses, which renders them highly vulnerable to the entry of pollutants and non-indigenous species to their ecosystem (Mirza *et al.*, 2014). Several studies have been conducted on the changes of diversity and number of planktons in the ballast tank of ships in

the Persian Gulf, such as plankton study of tanks on 2 oil tankers in Shahid Rajaei Port (Sangpour *et al.*, 2010; Shapoori and Gholami, 2014), and plankton study in the ROPME Sea Area (Dorgham, 2013), as well as other sea areas, including Malaysia (Siang *et al.*, 2018) and Cooper Port (Slovenia) (David *et al.*, 2007) with a focus on studying the impact of ballast water exchange on plankton population changes.

Shahid Rajaei Port is located on the entrance of the Persian Gulf, Strait of Hormuz, and international waterways, where various ships, container carriers, and oil tankers navigate, load, and discharge their cargo. The high traffic of such ships to this port is thus accompanied by the introduction of different non-indigenous and invasive species into the region through ballast water on ships. The issue of identifying threatening factors that enter the marine environment in this manner and result in reduced or lost efficiency and health of the ecosystems is therefore crucial and inevitable as an asset to satisfactory management, as the restoration of the resulting changes to many of such ecosystems varies between severely difficult and practically impossible. In line with the above necessity, the present study was conducted to identify and study the diversity and density of phytoplankton, as well as the possibility for transfer of phytoplankton species through the ballast water of ships entering Shahid Rajaei Port (Sangpour

et al., 2010; Shapoori and Gholami, 2014; Tolian *et al.*, 2020)

Materials and methods

Shahid Rajaei Port is located in Hormozgan Province, Iran (N: 27° 07'; E: 56° 04') (Shirowzhan *et al.*, 2016). The sampling for the present study was conducted on 24 ships entering this port from origins located in four regions, including the Persian Gulf (Iraq,

Kuwait, Qatar, and the United Arab Emirates), eastern Indian Ocean (Malaysia, Indonesia, and Singapore), western Indian Ocean (Kenya, Djibouti, and Somalia), and northwestern Indian Ocean (Sri Lanka, India, and Pakistan) (Fig. 1), in two seasons of winter 2017 and summer 2018. The particulars of the ships involved in the present study are presented in Table 1.




Figure 1: Geographical location of Shahid Rajaei Port, Hormozgan, Iran.

Sampling

The sampling was conducted in compliance with the G2 standard method approved by MEPC.173 (58) of

the International Maritime Organization (IMO), and during the berthing of ships at the mentioned port (MEPC, 2008).

Table1: Specifications of the 24 sampled ships in the Shahid Rajaei Port, Hormozgan, Iran.

Ships No	Ballast water Source	Ship Type	Gross Tonnage (t)	Season of sampling
S1	UMM QASR-IRAQ	General Cargo	34859	winter
S2	SHUWAIKH PORT- KUWAIT	Bulk Carrier	32983	winter
S3	HAMAD PORT- QATAR	Bulk Carrier	32945	winter
S4	JEBELALI-U.A.E	OIL Tanker	40037	summer
S5	MINA ZAYED- U.A.E	Bulk Carrier	17997	summer
S6	UMM QASR -IRAQ	Bulk Carrier	17979	summer
S7	MOMBASA- KENYA	Container	23389	winter
S8	MOMBASAKENYA	General Cargo	15670	winter
S9	MOMBASA- KENYA	General Cargo	16694	winter
S10	BERBERA- SOMALIA	Bulk Carrier	11743	summer
S11	DJIBOTI PORT	Bulk Carrier	33037	summer
S12	DJIBOTI PORT	Bulk Carrier	15609	summer
S13	PORT PASIR GUNAYNG - MALAYSIA	Bulk Carrier	33044	winter
S14	KENDAWANGAN- INDONESIA	General Cargo	25676	winter
S15	SINGAPORE PORT	General Cargo	36259	winter
S16	SINGAPORE PORT	Bulk Carrier	32305	summer
S17	SUNGAI PAKNING-INDONESIA	General Cargo	19354	summer
S18	SINGAPORE	OIL Tanker	29266	summer
S19	TRINCOMALEE- SRILANKA	OIL Tanker	32474	winter
S20	MAGDALLA- INDIA	OIL Tanker	32725	winter
S21	MUNDRA- INDIA	Bulk Carrier	32474	winter
S22	CHITTAGONG- BANGLADESH	Bulk Carrier	32305	summer
S23	KARACHI- PAKISTAN	Bulk Carrier	32945	summer
S24	GMB DAHEJ PORT- INDIA	Bulk Carrier	31117	summer

A total of 24 ships, including 13 bulk carriers, 4 oil tankers, 6 general cargo and 1 container, were sampled for phytoplankton existing in the ballast water. For 20 ships, the overflow method from the air pipe of the bottom tank was used, and on ships no. 1, 20, and 21 were sampled through the discharge valve near the discharge line. Considering the rapid loading and discharge of cargos ships, and in order not to delay the ships' operations, as well as safety considerations for the container carrier, the sampling on the container carrier was conducted by the crew through the discharge valve in the engine room.

For sampling purposes, 100 liters of ballast water was passed through a 20-micron plankton net, using a bucket, and the filtered samples were collected in a special container, fixed with 4-percent formalin, and transferred to the laboratory (David *et al.*, 2007; Siang *et al.*, 2018). Physical parameters inside ship tanks could affect the population of transferred species, and several such parameters, including temperature, salinity, and pH was thus

Phytoplankton Identification

To identify and study the density of phytoplankton at the species level, 3 takes (1 mL) of each sample were

transferred onto Sedgewick Rafter counting chambers and studied by an inverted microscope (Axiovert S100), using standard identification keys (Hasle *et al.*, 1996; Tomas, 1997; Al-Kandari *et al.*, 2009).

Statistical analysis

The collected data was initially assessed through the Kolmogorov-Smirnov Test to ensure their normalcy, and then significant difference among parameters for sampling stations and seasons was considered through the One-Way ANOVA. Average values were also compared through Duncan multiple range tests, and Pearson Correlation Coefficient among the number of species, abundance, and the measured water parameters was calculated using the SPSS V. 21. The seasonal similarity of the sampled phytoplankton was determined through PAST and Ecological Methodology software packages, while the cluster analysis and PCA test were performed by PRIMER Version 5 and PAST Version 2.13, respectively.

Results

Water parameters

The minimum and maximum ballast water temperature of 24 ships in winter 2017 and summer 2018 varied between 26.43° C and 40.06°C (average 33.24°C) in winter. In the cold season, salinity was between 34.76 and 38.33 ppt, the Persian Gulf had the highest registered level, while this parameter did not have any significant difference for the ballast water on ships loaded at the studied regions of the Indian Ocean ($p < 0.05$). In the same season, the pH for ballast water taken from ships entering the port from origins in the western Indian Ocean registered the highest average of 8.10 ($p > 0.05$). In the warm season, none of the measured parameters registered significant differences ($p < 0.05$) between the Persian Gulf and the three studied regions of the Indian Ocean. The average value of ballast water physical parameters measured in the summer was 7.96 for pH, 36.73 ppt for salinity, and 38.60°C for temperature (Table 2).

Table 2: Measured physical parameters of 24 ship's ballast water in winter 2017 and summer 2018, Hormozgan, Iran.

Location	Season	Temperature (°C)	Salinity(ppt)	pH
Persian Gulf	Winter	26.43±1.81 ^b	38.33±0.26 ^a	7.84±0.05 ^{ab}
	Summer	37.8±0.36 ^a	38.16±0.98 ^a	7.81±0.05 ^b
West Indian Ocean	Winter	28.36±0.16 ^b	35.46±0.4 ^{ab}	8.1±0.05 ^{ab}
	Summer	37.43±0.6 ^a	36.36±0.92 ^{ab}	7.76±0.23 ^b
East Indian Ocean	Winter	28.7±0.57 ^b	33.33±0.36 ^b	7.87±0.02 ^{ab}
	Summer	40.06±0.57 ^a	35.46±0.35 ^{ab}	8.23±0.07 ^a
Northwest Indian Ocean	Winter	28.06±1.66 ^b	34.76±1.61 ^{ab}	7.95±0.03 ^{ab}
	Summer	39.13±1.09 ^a	33.5±1.84 ^b	8.04±0.05 ^{ab}

The same letters mean no difference and different letters mean a significant difference at the 5% level between experimental regions ($p < 0.05$).

The values for ballast water temperature, salinity, and pH on ships entering Shahid Rajaei Port in winter 2017 and summer 2018 are depicted in Fig. 2. The values included limited fluctuation, and the only temperature

registered a significant difference between summer and winter ($p>0.05$), meaning that on all ships understudy, the water temperature was higher in summer than winter ($p<0.05$).




Figure 2: Measured physical parameters of 24 ship's ballast water entered to Shahid Rajaei Port in winter 2017 and summer 2018, Hormozgan, Iran.

Phytoplankton population composition

The results of sampling phytoplankton found in the ballast water of ships under study are presented in Table 3. The species identified in this study belonged to 4 phyla, 5 classes, 23 orders, 38 families, and 108 species. After studying the ballast tanks, it was determined that classes of Dinophyceae (with 6 orders, 14 families and 39 species) and Bacillariophyceae (with 13 orders, 18 families, and 57 species)

accounted for 48 and 46 percent of the species respectively, while the classes of Raphidophyceae and Cyanophyceae (with 1 order, 1 family and 1 species) composed only 1 percent of the species identified. Moreover, in the warm season, class Dinophyceae (54 percent), and in the cold season, class Bacillariophyceae (60.96 percent) were prevalent among the samples (Figs. 3 and 4).

Table 3: Classification of phytoplankton observed in ballast water tanks on ships entering Shahid Rajaei Port (winter 2017 & summer 2018).

Phylum	Class	Order	Family	Genus/species	HAB
Myzozoa	Dinophyceae	Gonyaulacales	Ceratiaceae	<i>Ceratium breve</i>	
				<i>Ceratium macroceros</i>	
				<i>Ceratium dens</i>	
				<i>Ceratium furca</i>	Al- Kandari <i>et al.</i> , 2009
				<i>Ceratium fusus</i>	Al- Kandari <i>et al.</i> , 2009
				<i>Ceratium trichoceros</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Ceratium lineatum</i>	
				<i>Ceratium massiliense</i>	
				<i>Ceratium tripos</i>	
				<i>Ceratium symmetricum</i>	
			Ostreopsidaceae	<i>Alexandrium minutum</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Chaetoceros affinis</i>	
				<i>Chaetoceros curvisetus</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Chaetoceros denticulatus</i>	
				<i>Chaetoceros socialis</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Chaetoceros subtilis</i>	
				<i>Chaetoceros peruvianus</i>	Al-Yamani <i>et al.</i> , 2019(II) Al-Kandari <i>et al.</i> , 2009

Table 3 (Continued):

Phylum	Class	Order	Family	Genus/species	HAB
			Gymnodiniaceae	<i>Cochlodinium polykrikoides</i> <i>Gonyaulax polygramma</i>	SMR Fatemi <i>et al.</i> , 2012; Matsuoka <i>et al.</i> , 2010 Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009; Saraji, 2018
				<i>Gyrodinium fusiforme</i> <i>Gyrodinium spirale</i> <i>Gymnodinium catenatum</i>	Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009; Saraji, 2018
			Polykrikaceae	<i>Polykrikos kofoidii</i>	
			Pyrophacaceae	<i>Pyrophacus steinii</i> <i>Pyrophacus horrologium</i>	
			Peridiniaceae	<i>Peridinium quinquecorne</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Scrippsiella trochoidea</i>	Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009
		Dinophysiales	Dinophysaceae	<i>Dinophysis rotundata</i> <i>Dinophysis acuta</i> <i>Dinophysis miles Cleve</i> <i>Dinophysis caudata</i>	Al-Yamani <i>et al.</i> , 2019(I) Al-Kandari <i>et al.</i> , 2009 Al-Yamani <i>et al.</i> , 2019(I) Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009
		Peridiniales	Oxytoxaceae	<i>Ornithocercus magnificus</i> <i>Oxytoxum scolopax Stein</i>	
			Protoperidiniaceae	<i>Protoperidinium claudicans</i> <i>Protoperidinium oceanicum</i> <i>Protoperidinium pyriforme</i> <i>Protoperidinium elegans</i>	

Table 3 (Continued):

Phylum	Class	Order	Family	Genus/spices	HAB	
Ochrophyta	Bacillariophyceae	Prorocentrales	Brachidiniaceae	<i>Protoperidinium pellucidum</i>	Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009	
				<i>karenia cf. brevis</i>		
			Goniodomataceae	<i>Pyrodinium bahamense</i>	Al-Yamani <i>et al.</i> , 2019(I); Al-Kandari <i>et al.</i> , 2009 Saraji, 2018	
			Prorocentrales	Prorocentraceae	<i>Prorocentrum gracile</i>	Al-Yamani <i>et al.</i> , 2019(I); Al- Kandari <i>et al.</i> , 2009
					<i>Prorocentrum minimum</i>	Al- yamani <i>et al.</i> , 2019(I)
					<i>Prorocentrummicans</i>	Al-Yamani <i>et al.</i> , 2019(I)
					<i>Prorocentrum balticum</i>	Al-Yamani <i>et al.</i> , 2019(I) Al- Kandari <i>et al.</i> , 2009
			Thoracosphaerales	Thoracosphaeraceae	<i>Goniodoma polyedricum</i>	
			Pyrocystales	Pyrophacaceae	<i>Pyrocystis noctiluca</i>	
			Chaetocerotana eincertaesedis	Chaetocerotaceae	<i>Chaetoceros pseudocurvisetum</i>	Al-Yamani <i>et al.</i> , 2019(I)
		<i>Bacteriastrumfurcatum</i>			Al-Yamani <i>et al.</i> , 2019(II)	
		Lithodesmiales	Lithodesmiaceae	<i>Bacteriastrum hyalinum</i>		
				<i>Helicotheca tamesis</i>		
		Hemiaulales	Bellerocheaceae	<i>Ditylum brightwellii</i>		
				<i>Bellerocheahorologicalis</i>		
		Hemiaulales	Bellerocheaceae	<i>Bellerochea malleus</i>		
				<i>Climacodium frauenfeldianum</i>		
<i>Eucampia zoodiacus</i>						
Hemiaulales	Hemiaulaceae	<i>Hemidiscus cuneiformis</i>				
		<i>Eucampia</i> sp.				
		<i>Coscinodiscus wailesii</i>				
Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus</i>				
		<i>asteromphalus</i>				

Table 3 (Continued):

Phylum	Class	Order	Family	<i>nb</i>	HAB
				<i>Coscinodiscus jonesianus</i>	
				<i>Coscinodiscus granii</i>	
				<i>Coscinodiscus marginatus</i>	
				<i>Coscinodiscus radiatus</i>	
				<i>Palmeria hardmaniana</i>	
		Bacillariales	Bacillariaceae	<i>Bacillaria paxillifera</i>	
				<i>Bacillaria socialis</i>	
				<i>Cylindrotheca closterium</i>	Al-Yamani <i>et al.</i> , 2019(II)
					Al-Kandari <i>et al.</i> , 2009
				<i>Nitzschia longissima</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Nitzschia behrei</i>	
				<i>Nitzschia sigma</i>	
				<i>Pseudonitzschia delicatissima</i> group	
				<i>Pseudonitzschia seriata</i> group	
		Triceratiales	Triceratiaceae	<i>Odontella longicuris</i>	
				<i>Odontella mobiliensis</i>	
		Asterolamprales	Asterolampraceae	<i>Asteromphalus ehrenberg</i>	
				<i>Asteromphalus flabellatus</i>	
		Thalassiosirales	Stephanodiscaceae	<i>Cyclotella</i> sp.	
				<i>Cyclotella stylorum</i>	
				<i>Brightwell</i>	
			Thalassiosiraceae	<i>Planktoniella sol</i>	
				<i>Thalassiothrix frauenfeldii</i>	
				<i>Thalassiosira eccentrica</i>	
			Lauderiaceae	<i>Lauderia annulata</i>	
		Naviculales	Diploneidaceae	<i>Diploneis splendida</i>	
		Leptocylindrales	Naviculaceae	<i>Navicula directa</i>	
				<i>Navicula elegans</i>	

Table 3 (Continued):

Phylum	Class	Order	Family	Genus/species	HAB
				<i>Navicula</i> sp.	
			Pleurosigmataceae	<i>Pleurosigma normanii</i>	
				<i>Pleurosigma elongatum</i>	
				<i>Pleurosigma</i>	
				<i>planctonicum</i>	
				<i>Pleurosigma diverse-</i>	
				<i>striatum</i>	
				<i>Pleurosigma formosum</i>	
				<i>Pleurosigma angulatum</i>	
			Pleurosigmataceae	<i>Pleurosigma strigosum</i>	
			Leptocylindraceae	<i>Leptocylindrus minimus</i>	
		Rhizosoleniaceae	Rhizosoleniaceae	<i>Dactyliosolen</i>	
				<i>fragilissimus</i>	
				<i>Guinardia flaccida</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>Guinardia striata</i>	
				<i>Proboscia alata</i>	
				<i>Rhizosolenia bergonii</i>	
			Rhaphoneidaceae	<i>Diplomenora</i>	
				<i>cocconeiforma</i>	
				<i>Rhizosolenia robusta</i>	
		Mastogloiales	Mastogloiaceae	<i>Mastogloia arabica</i>	
		Melosirales	Stephanopyxidaceae	<i>Stephano</i>	
				<i>pyxispalmeriana</i>	
	Dictyochophyceae	Dictyochales	Dictyochaceae	<i>Dictyocha fibula</i>	Al- Kandari <i>et al.</i> , 2009
		Surirellales	Surirellaceae	<i>Surirella splendida</i>	
				<i>Surirella fastuosa</i>	
Chromophyta	Raphidophyceae	Chattonellales	Chattonellaceae	<i>Heterosigma</i> sp.	
Cyanobacteria	Cyanophyceae	Oscillatoriales	Phormidiaceae	<i>Trichodesmium</i>	Al-Yamani <i>et al.</i> , 2019(I)
				<i>erythraeum</i>	Al- Kandari <i>et al.</i> , 2009

According to Fig. 5, Bacillariophyceae was the most abundant class found on ballast waters from the Persian Gulf and western Indian Ocean, and Dinophyceae on ships from the eastern and northwestern Indian Ocean. The density of phytoplankton also depicted that ships from the western

Indian Ocean carried the highest density of phytoplankton in winter and ships from the eastern Indian Ocean in summer. The lowest density of ballast water phytoplankton was observed on ships from the northwestern Indian Ocean (Fig. 6).




Figure 3: Abundance percentage (%) of identified prevalent phytoplankton; A: Total study period; B: winter 2017; C: summer 2018) in the ballast water of entered ships to Shahid Rajaei Port, Hormozgan, Iran.




Figure 4: Abundance percentage (%) of identified prevalent phytoplankton species in the ballast water of entered ships (A: in winter 2017; B: in summer 2018) to Shahid Rajaei Port, Hormozgan, Iran.




Figure 5: Main observed phytoplankton classes by regional origin in the ballast water of entered ships to Shahid Rajaei Port, Hormozgan, Iran.




Figure 6: Impact of regional origin on the density of phytoplankton (Cells per Liter) in the ballast water of entered ships to Shahid Rajaei Port, Hormozgan, Iran.

Ceratium breve and *Proboscia alata* were present in all regions in two seasons except on ships from the northwest Indian Ocean in winter. Different phytoplankton species are presented in Table 4 by season and region.

The results of the Pearson Correlation and PCA Test pointed to a positive correlation at 0.05 percent between phytoplankton abundance and temperature. In other words, the number of phytoplankton abundance increased with the increase in temperature of ballast water (Table 5 and Fig. 7).

Cluster analysis of similarity among regional origins of ships entering Shahid Rajaei Port

The results of the similarity test following Jaccard and Sorensen Indexes showed the highest similarity to exist between WW and WS (Jaccard: 0.563; Sorensen 0.72), and the lowest similarity between PGS and NW (Jaccard: 0.111; Sorensen 0.2), as presented in Table 6.

According to the above cluster analysis and identified phytoplankton, ships arriving from the western Indian Ocean had the highest similarity of phytoplankton density averages (64 percent) during winter and summer,

while ships from the eastern and northwestern Indian Ocean registered the lowest similarity (44 and 4 percent,

respectively) during winter and summer, with other regions under study (Fig. 8).

Table 4: Seasonal comparison of phytoplankton presence in ballast water tanks on ships entering Shahid Rajaei Port (winter 2017 & summer 2018), Hormozgan, Iran.

Species	Indian Ocean				Indian Ocean			
	Persian Gulf	West	East	North west	Persian Gulf	West	East	North west
<i>Alexandrium minutum</i>		+	+			+	+	
<i>Asteromphalus ehrenberg</i>		+					+	
<i>Asteromphalus flabellatus</i>		+				+		
<i>Bacillaria paxillifera</i>	+							
<i>Bacillaria socialis</i>							+	
<i>Bacteriastrium furcatum</i>		+	+					+
<i>Bacteriastrium hyalinum</i>			+				+	
<i>Bellerochea horologicalis</i>		+			+	+		
<i>Bellerochea malleus</i>		+	+				+	
<i>Ceratium breve</i>	+	+	+		+	+	+	+
<i>Ceratium dens</i>		+						
<i>Ceratium furca</i>	+	+	+	+		+	+	+
<i>Ceratium fusus</i>			+		+		+	+
<i>Ceratiumlineatum</i>		+	+			+	+	
<i>Ceratium macroceros</i>			+		+		+	
<i>Ceratium massiliense</i>		+	+			+	+	+
<i>Ceratium symmetricum</i>	+	+				+	+	
<i>Ceratium trichoceros</i>				+	+			+
<i>Ceratium tripos</i>					+	+		+
<i>Chaetoceros affinis</i>					+			
<i>Chaetoceros curvisetus</i>						+		
<i>Chaetoceros peruvianus</i>		+				+	+	+
<i>Chaetoceros pseudocurvisetum</i>		+			+			
<i>Chaetoceros socialis</i>			+			+		
<i>Chaetoceros subtilis</i>		+				+	+	+
<i>Climacodium frauenfeldianum</i>				+				
<i>Cochlodinium polykrikoides</i>							+	
<i>Coscinodiscus asteromphalus</i>		+					+	
<i>Coscinodiscus granii</i>	+			+	+	+	+	
<i>Coscinodiscus jonesianus</i>							+	
<i>Coscinodiscus marginatus</i>			+	+				
<i>Coscinodiscus radiatus</i>	+		+	+				
<i>Coscinodiscus wailesii</i>							+	+
<i>Cyclotella</i> sp.			+				+	+
<i>Cyclotella stylorum</i> Brightwell	+	+				+	+	
<i>Cylindrotheca closterium</i>			+			+	+	
<i>Dactyliosolen fragilissimus</i>							+	
<i>Dictyocha fibula</i>	+					+		+
<i>Dinophysis acuta</i>		+					+	+
<i>Dinophysis caudata</i>	+	+	+	+	+	+	+	+
<i>Dinophysis cleve miles</i>	+							
<i>Dinoysis rotundata</i>	+				+		+	

Table 4 (Continued):

Species	Indian Ocean				Indian Ocean			
	Persian Gulf	West	East	North west	Persian Gulf	West	East	North west
	Winter				Summer			
<i>Diplomenora cocconeiforma</i>			+				+	
<i>Diploneis splendida</i>				+				
<i>Ditylum brightwellii</i>						+	+	
<i>Eucampia</i> sp.						+		
<i>Eucampia zoodiacus</i>	+	+		+		+	+	
<i>Goniodomapedricum</i>							+	
<i>Gonyaulax polygramma</i>				+			+	
<i>Guinardia flaccida</i>	+	+		+				+
<i>Guinardia striata</i>	+						+	
<i>Gymnodinium catenatum</i>					+			
<i>Gyrodinium fusiforme</i>					+		+	
<i>Gyrodinium spirale</i>					+			
<i>Helicotheca tamesis</i>				+				
<i>Hemidiscus cuneiformis</i>								+
<i>Heterosigma</i> sp.			+		+		+	
<i>karenia cf. brevis</i>							+	+
<i>Lauderia annulata</i>					+		+	
<i>Leptocylindrus minimus</i>			+					
<i>Mastogloia arabica</i>		+				+	+	
<i>Navicula directa</i>			+	+				
<i>Navicula elegans</i>		+			+			+
<i>Navicula</i> sp.			+				+	
<i>Nitzschia behrei</i>							+	+
<i>Nitzschia longissima</i>			+					+
<i>Nitzschia sigma</i>		+			+	+		+
<i>Odontella longicuris</i>					+		+	
<i>Odontella mobiliensis</i>		+				+		
<i>Ornithocercus magnificus</i>	+	+				+		+
<i>Oxytoxum scolopax</i> Stein	+							
<i>Palmeria hardmaniana</i>	+							
<i>Peridinium quinquecorne</i>	+					+		+
<i>Planktoniella sol</i>	+	+				+		+
<i>Pleurosigma angulatum</i>			+					+
<i>Pleurosigma diverse-striatum</i>			+			+	+	
<i>Pleurosigma elongatum</i>	+	+				+		
<i>Pleurosigma formosum</i>			+					
<i>Pleurosigma normanii</i>				+			+	
<i>Pleurosigma planctonicum</i>							+	
<i>Pleurosigma strigosum</i>							+	
<i>Polykrikos kofoidii</i>					+	+		
<i>Proboscia alata</i>	+	+	+		+	+	+	+
<i>Prorocentrum balticum</i>			+				+	+
<i>Prorocentrum gracile</i>	+	+		+	+	+		+
<i>Prorocentrummicans</i>	+	+			+	+		+
<i>prorocentrum minimum</i>		+						+
<i>Protoperidinium claudicans</i>		+				+		
<i>Protoperidinium oceanicum</i>								+
<i>Protoperidinium pellucidum</i>		+					+	
<i>Protoperidinium pyriforme</i>		+				+		

Table 4 (Continued):

Species	Indian Ocean				Indian Ocean			
	Persian Gulf	West	East	North west	Persian Gulf	West	East	North west
	Winter				Summer			
<i>Protopheridinium elegans</i>								+
<i>Pseudonitzschia delicatissima</i> group	+	+				+	+	
<i>Pseudonitzschia seriata</i> group			+	+			+	
<i>Pyrocystis noctiluca</i>							+	
<i>Pyrodinium bahamense</i>		+	+			+	+	+
<i>Pyrophacus horroloogium</i>			+					+
<i>Pyrophacus steinii</i>								+
<i>Rhizosolenia bergonii</i>		+		+		+		
<i>Rhizosolenia robusta</i>				+				
<i>Scrippsiella trochoidea</i>		+				+		
<i>Stephanopyxis palmeriana</i>							+	
<i>Surirella fastuosa</i>				+				
<i>Surirella splendida</i>			+					+
<i>Thalassiosira eccentrica</i>	+	+			+	+		+
<i>Thalassiothrix frauenfeldii</i>			+		+		+	
<i>Trichodesmium erythraeum</i>	+		+		+	+	+	

Table 5: Correlation between measured parameters and phytoplankton density in the ballast water tanks on entered ships to Shahid Rajaee Port, Hormozgan, Iran.

	Phytoplankton Correlation	Taxa	Abundance
Temperature	Pearson Correlation	.389	.483*
	Sig. (2-tailed)	.060	.017
	N	24	24
Salinity	Pearson Correlation	-.075	-.054
	Sig. (2-tailed)	.726	.801
	N	24	24
pH	Pearson Correlation	.080	-.029
	Sig. (2-tailed)	.712	.894
	N	24	24




Figure 7: Principle Component Analysis (PCA) to consider the correlation between ballast water parameters and phytoplankton density in the ballast water tanks of entered ships to Shahid Rajaee Port, Hormozgan, Iran.

Table 6: Similarity among Regional Origins in the ballast water tanks of entered ships to Shahid Rajaei Port, Hormozgan, Iran.

Region and Season	Index	PGW*	PGS*	WW*	WS*	EW*	ES*	NW*	NS*
PGW	Jaccard	1	0.178	0.28	0.37	0.113	0.196	0.195	0.326
	Sorenson	1	0.302	0.438	0.54	0.203	0.328	0.327	0.492
PGS	Jaccard		1	0.138	0.212	0.176	0.197	0.111	0.22
	Sorenson		1	0.242	0.349	0.3	0.329	0.2	0.361
WW	Jaccard			1	0.563	0.203	0.253	0.173	0.298
	Sorenson			1	0.72	0.338	0.404	0.295	0.459
WS	Jaccard				1	0.186	0.227	0.132	0.309
	Sorenson				1	0.314	0.37	0.233	0.472
EW	Jaccard					1	0.338	0.143	0.193
	Sorenson					1	0.506	0.25	0.324
ES	Jaccard						1	0.147	0.2
	Sorenson						1	0.256	0.333
NW	Jaccard							1	0.118
	Sorenson							1	0.211
NS	Jaccard								1
	Sorenson								1

*PG: Persian Gulf, W: west Indian Ocean, E: east Indian Ocean, N: North west Indian Ocean, W: Winter 2017, S: Summer 2018.




Figure 8: Similarity cluster analysis among regional origins in the ballast water tanks of entered ships to Shahid Rajaei Port, Hormozgan, Iran.

PG: Persian Gulf, W: West Indian Ocean, E: East Indian Ocean, N: North west Indian Ocean, W: Winter 2017, S: Summer 2018.

Discussion

Ballast water carried on ships is a way which species are transferred from one aquatic environment to another and the consequences of such transfer exceed the mere introduction of pollutants such

as oil, heavy metals, and persistent organic pollutants. Invasive and non-indigenous species either die under new environmental conditions or develop and spread, which leaves initially intangible and small impacts that

aggravate and accelerate through time (Steichen *et al.*, 2012; Vieira *et al.*, 2018).

Uncontrolled exchange of ballast water and entry of invasive aquatic species is a growing concern that results in ecologic, economic, and health-related damages to the host environment (Werschkun *et al.*, 2014; Wu *et al.*, 2017). Hence, minimizing such invasion and transfer through ballast water carried on ships can be identified as a major priority for the maritime industries around the world (Balaji *et al.*, 2014; de Castro *et al.*, 2017).

Considering the importance of ballast water and the mandatory requirements to inspect it under articles 6, 9 and 10 of the convention focused on monitoring and promoting ballast water management in Iran (Bobka, 2018; Rey *et al.*, 2018). The present study proceeded to analyze ballast water on board 24 ships, (including 13 bulk carriers, 4 oil tankers, 6 general cargo and 1 container) entering Shahid Rajaei Port in terms of phytoplankton species present, and the impact of ballast water physical parameters on their survival.

Changes in the physical parameters of the environment affect the survival of aquatic species to different degrees. In the present study, temperature as the only effective parameter on phytoplankton composition did not significantly differ among regional origins of ships (26.43°C to 28.70°C in winter, and 37.43°C to 40.06°C in summer), or with the temperature range

in the Persian Gulf (32° C to 34°C) (Fazelpoor *et al.*, 2016) (Table 5 and Fig. 7). With its impact on plankton metabolism, growth, enzyme activity, and cellular osmosis stress (D'ors, 2016), the temperature is considered among the most decisive parameters affecting micro-organic survival in ballast water tanks on ships.

The absence of significant temperature difference between origin and destination of ships thus confirms the similarity of phytoplankton species in such regions, and extensively increases the possibility of survival for species transferred from those regional origins. In a study by Siang *et al.* (2018), temperature and salinity were introduced as parameters affecting phytoplankton survival, and the impact of small temperature differences among ports of origin and destination on micro-organic survival after transfer into the new environment through ballast water discharge was highlighted, which corroborates the findings of the present study.

It was also observed that the number of phytoplankton in ballast water tanks was significantly higher in summer than winter ($p < 0.05$), in a manner, that the average number of phytoplankton present in the ballast water of ships arriving from the Persian Gulf, western, eastern, and the northwestern Indian Ocean was measured as 655.65, 944.42, 1383.55 and 625.97 species in 100-liter water, respectively, in summer, and 237.72, 575.40, 908.05, and 159.62 species in 100-liter water, respectively,

in winter. The reason can be attributed to the occurrence of stronger monsoons in summer than winter. The seasonal winds disrupt the water column and seabed structure, resulting in the higher dissolution of oxygen and other materials in water and the creation of more favorable living conditions for phytoplankton (Mitbavkar and Anil, 2006).

The ballast water tanks on all 24 ships entering Shahid Rajaei Port (the Persian Gulf) contained phytoplankton species: ships arriving from the Persian Gulf 43 species, western Indian Ocean 52 species, eastern Indian Ocean 65 species, and northwestern Indian Ocean 51 species. The results thus showed that incoming ships from the eastern Indian Ocean carried the highest number and density of phytoplankton in their tanks. Furthermore, the analysis of species and regional origins of the ships pointed to the similarity between species identified on ships from the Persian Gulf and western Indian Ocean (Fig. 8). The reason is the prevalent conditions in the Gulf of Oman at the connection of that Gulf to the Indian Ocean, the westbound currents in the northern parts, and eastbound currents on the southern coasts, which guide the water of the Indian Ocean (particularly in the west) toward the northern Gulf of Oman, and into the Persian Gulf through the Strait of Hormuz (Layeghi *et al.*, 2019), and resulting in the highly similar biological conditions and plankton composition in the Persian Gulf and western Indian Ocean. The

prevalence of class Bacillariophyceae in the ballast water further confirms such similarity, which can be attributed to the existence of silicate in ballast sediments as a nutrient for diatom growth and higher density and diversity of the mentioned class (Akbarzadeh *et al.*, 2017).

The lowest density and number of samples and similarity between species was observed in the water coming from the Northwestern Indian Ocean, possibly due to the treatment system of the ballast water on ships no. 20 and 21 that arrived in Shahid Rajaei Port from India. The class of diatoms or Bacillariophyceae from the Ochrophyta phylum was frequent in the winter, because of the cryophilic behavior of the mentioned species (Hegedus *et al.*, 2009). With the reduced frequency of this class under higher temperatures in summer, the class of Dinophyceae from the Myzozoa phylum replaced the Bacillariophyceae as the prevalent class.

Among the identified species, *Dinophysis clevevilles*, *Oxytoxum scolopax Stein*, *Palmeria hardmaniana*, and *Bacillaria paxillifera* were only observed in ballast water from the Persian Gulf, *Ceratium dens*, *Chaetoceros curvisetus*, *Scrippsiella trochoidea*, *Protoperidinium claudican*, *Protoperidinium pyriforme*, *Eucampia* sp., *Odontella mobiliensis*, and *Asteromphalus flabellatus* from the western Indian Ocean, *Cochlodinium polykrikoides*, *Goniodoma polyedricum*, *Pyrocystis noctiluca*, *Bacteriastrum*

hyaline, *Coscinodiscus jonesianus*, *Bacillaria socialis*, *Navicula* sp., *Pleurosigma planctonicum*, *Pleurosigma formosum*, *Pleurosigma strigosum*, *Leptocylindrus minimus*, *Diplomenora cocconeiforma*, and *Stephanopyxi spalmeriana* only from the eastern Indian Ocean, and *Helicotheca tamesis*, *Climacodium frauenfeldianum*, *Diploneis splendida*, *Rhizosolenia robusta*, *Surirella fastuosa*, *Hemidiscus cuneiformis*, *Protoperidinium elegans*, and *Pyrophacus steinii* in the ballast water from the northwestern Indian Ocean. The species common between the Indian Ocean and the Persian Gulf were *Ceratium breve*, *Ceratium furca*, *Dinophysis caudate*, and *Proboscia alata*. Dinoflagellates are harmful species that result in red tide in the region. The class of Dinophysiales accounted for 3.88 percent of the identified species on average, with 4.05 percent in summer and 3.54 percent in winter. The harmful species identified in this study included *Dinophysis rotundata*, *Dinophysis acuta*, *Dinophysis miles* Cleve, *Dinophysis caudate*, *Peridinium quinquecorne*, and *Gonyaulax polygramma*, which not only pollutes marine species but also jeopardizes human life through intoxication. The presence of such invasive and dangerous species have been previously reported in studies by Lim *et al.* (2004; 2012) and Teen *et al.* (2012) in the ballast water of ships entering in Kuantan Port of Malaysia, Roziawati *et al.* (2015) for Malacca,

Tan *et al.* (2016) for Singapore, and Siang *et al.* (2018) for Malaysia. Out of the 108 species identified in the present research, 28 species were reported for causing red tide, such as *Prorocentrum gracile* in Qatar, *Prorocentrum balticum*, *Prorocentrum minimum*, *Peridinium quinquecorne*, *Guinardia flaccida*, *Prorocentrum balticum*, *Bacteriastrum furcatum*, *Gonyaulax polygramma* in Oman, *Trichodesmium erythraeum* in Oman and the Arabian Sea, *Cylindrotheca closterium* and *bahamense* in the United Arab Emirates. Others have caused death among animals and humans, such as *Alexandrium minutum*, *Ceratium furca*, *Ceratium fusus*, *prorocentrum minimum*, *Scrippsiella trochoidea*, and *Prorocentrum micans* (Table 3).

Thalassiosira and *Chaetoceros*, known for surviving extensive light levels (Liebich *et al.*, 2012) and causing extensive and harmful red tide (Shen *et al.*, 2012), were also observed among the Phytoplankton identified by the present study at Shahid Rajaei Port (Sidabutar *et al.*, 2021). Among the 108 species identified by the present study in the ballast water of incoming ships, *Odontella longicruris* was the only species not previously identified and reported by relevant studies in the Persian Gulf (Al-Kandari *et al.*, 2009; Saraji, 2018; Al-Yamani *et al.*, 2019). Based on the findings of the present study, although invasive or non-indigenous species were not observed among the identified species, and apart from the *Odontella longicruris* not

previously reported in the region, all other species belonged to the phytoplankton composition of the host ecosystem, yet their unnatural presence in the new environment and survival capability due to negligible difference in temperature could be deemed as possible conditions for growing population and prevalence of a particular species, and the resulting adverse consequences, such as red tide and due to the high traffic of ships through the Strait of Hormuz and the possible transfer of invasive species to this port, however, the survival of even one non-indigenous species in the new environment could culminate in irreparable damages if not properly controlled and managed.

Acknowledgment

This paper was published under the valuable cooperation of the Port State Control Officer at Shahid Rajaei Port, who lent their assistance in sampling ballast water on ships, and with the financial and spiritual support of the Iranian Ports and Maritime Organization.

References

- Akbarzadeh, G.H., Salarpoori, A., Darvishi, M. and Behzadi, S., 2017.** Spatial and temporal variations of phytoplankton population in coastal waters of Hormozgan province (Qeshm and leangeh). *Iranian Scientific Fisheries Journal*, 25(5), 11-21. DOI: 10.1016/j.hal.2006.11.006
- Al-Kandari, M., Al-Yamani, F.Y. and Al-Rifaie, K., 2009.** Marine phytoplankton atlas of Kuwait's waters. *Kuwait Institute for Scientific Research*, 351-366. DOI: 10.1016/j.hal.2006.11.006
- Al-Majed, N., Mohammadi, H., Al-Ghadban, A. and Al-Awadi, A.R., 2000.** *Regional report of the state of the marine environment*. Regional Organization for the Protection of the Environment.
- Al-Yamani, F. and Saburova, M.A., 2019.** Marine phytoplankton of Kuwait's waters. Volume II Diatoms. Cyanobacteria, Dinoflagellates, Flagellates. Kuwait.
- Balaji, R., Yaakob, O. and Koh, K.K., 2014.** A review of developments in ballast water management. *Environmental Reviews*, 22(3), 298-310. DOI: 10.1139/er-2013-0073
- Balley, S.A. and Munawar, M., 2015.** A synthesis of marine invasive species research and management in the ROPME sea area. *Aquatic Ecosystem Health and Management*, 18(3), 374-354. DOI: 10.1080/14634988.2015.1039917
- Bobka, J.R., 2018.** Defining Unduly: Resolving Inherent Textual Ambiguity in the IMO's Ballast Water Management Convention. *Wm. and Mary L. Rev.*, 60, 267P.
- Burkholder, J.M., Hallegraeff, G.M., Melia, G., Cohen, A., Bowers, H.A., Oldach, D.W. and Mallin, M.A., 2007.** Phytoplankton and

- bacterial assemblages in ballast water of US military ships as a function of port of origin, voyage time, and ocean exchange practices. *Harmful Algae*, 6(4),486-518. DOI: 10.1016/j.hal.2006.11.006
- Čampara, L., Slišković, M. and JelićMrčelić, G., 2019.** Key ballast water management regulations with a view on ballast water management systems type approval process. *NAŠE MORE: znanstveničasopisza more ipomorstvo*, 66(2),78-86. DOI 10.17818/NM/2019/2.5
- Cordell, J., Kalata, O., Pleus, A., Newsom, A., Strieck, K. and Gertsen, G., 2015.** Effectiveness of ballast water exchange in protecting Puget Sound from invasive species. Washington Department of Fish and Wildlife, pp. 1- 55.
- David, M., Gollasch, S., Cabrini, M., Perkovič, M., Bošnjak, D. and Virgilio, D., 2007.** Results from the first ballast water sampling study in the Mediterranean Sea—the Port of Koper study. *Marine Pollution Bulletin*, 54(1),53-65. DOI: 10.1016/j.marpolbul.2006.08.041
- De Castro, M.C.T., Fileman, T.W. and Hall-Spencer, J.M., 2017.** Invasive species in the Northeastern and Southwestern Atlantic Ocean: A review. *Marine Pollution Bulletin*, 116(1-2), 41-47. DOI: 10.1016/j.marpolbul.2016.12.048
- Dickman, M. and Zhang, F., 1999.** Mid-ocean exchange of container vessel ballast water. Effects of vessel type in the transport of diatoms and dinoflagellates from Manzanillo, Mexico, to Hong Kong, China. *Marine Ecology Progress Series*, 176, 253-262. DOI: 10.1016/j.marpolbul.2004.01.016
- Dorgham, M.M., 2013.** Plankton research in the ROPME sea area, achievements, and gaps. DOI: 10.22059/IJER 2013.656
- D'ors, A., Bartolomé, M.C. and Sánchez-Fortún, S., 2016.** Repercussions of salinity changes and osmotic stress in marine phytoplankton species. *Estuarine, Coastal and Shelf Science*, 175, 169-175. DOI:10.1016/j.ecss.2016.04.004
- Drake, J.M. and Lodge, D.M., 2004.** Global hot spots of biological invasions: evaluating options for ballast water management. Proceedings of the Royal Society of London. Series B: *Biological Sciences*, 271(1539), 575-580. DOI: 10.1098/rspb.2003.2629
- Drake, L.A., Doblin, M.A. and Dobbs, F.C., 2007.** Potential microbial bioinvasions via ships' ballast water, sediment, and biofilm. *Marine Pollution Bulletin*, 55(7-9), 333-341. DOI: 10.1016/j.marpolbul.2006.11.007
- Endresen, Ø., Behrens, H.L., Brynstad, S., Andersen, A.B. and Skjong, R., 2004.** Challenges in global ballast water management. *Marine Pollution Bulletin*, 48(7-8),615-623. DOI: 10.1016/j.marpolbul.2004.01.016

- Fatemi, S.M.R., Nabavi, S.M.B., Vosoghi, G., Fallahi, M. and Mohammadi, M., 2012.**The relation between environmental parameters of Hormuzgan coastline in Persian Gulf and occurrence of the first harmful algal bloom of *Cochlodinium polykrikoides* (Gymnodiniaceae). *Iranian Journal of Fisheries Sciences*, 11(3), 475-489. URL: <http://jifro.ir/article-1-621-en.html>
- Fazelpoor, K., DadollahiSohrab, A., Elmizadeh, H., Asgari, H.M. and Khazaei, SH., 2016.**The evaluation of sea surface temperature and the relationship between SST and depth in the Persian Gulf by MODIS.
- Gerhard, W.A., Lundgreen, K., Drillet, G., Baumler, R., Holbech, H. and Gunsch, C.K., 2019.** Installation and use of ballast water treatment systems—Implications for compliance and enforcement. *Ocean and Coastal Management*, 181, 104907. DOI: 10.1016/j.ocecoaman.2019.104907
- Gollash, S., David, M., Dragsund, E., Hewitt, C. and Fukuyo, Y., 2007.** Critical review of the IMO international convention on the management of ships' ballast water and sediment. *Harmful Alga*, 6(4), 585-600. DOI: 10.1016/j.hal.2006.12.009
- Hallegraeff, G.M. and Bolch, C.J., 1992.** Transport of diatom and dinoflagellate resting spores in ships' ballast water: implications for plankton biogeography and aquaculture. *Journal of Plankton Research*, 14(8), 1067-1084. DOI: 10.1093/plankt/14.8.1067
- Hasle, G.R., Syvertsen, E.E., Steidinger, K.A., Tangen, K. and Tomas, C.R., 1996.** Identifying marine diatoms and dinoflagellates. Elsevier. DOI: 10.1016/B978-0-12-693015-3.X5000-1
- Hegedus, C., Popescu, S., Boaru, A., Borda, C. and Fornade, A., 2009.** The Study of some Physical-Chemical Parameters of Water and Relationship with Phytoplankton in Fish Pond. *Bulletin UASVM Animal Science and Biotechnologies*, 66, 1-2.
- Lakshmi, E., Priya, M. and Achari, V.S., 2021.** An overview on the treatment of ballast water in ships. *Ocean and Coastal Management*, 199, 105296. DOI: 10.1016/j.ocecoaman.2020.105296
- Layeghi, B., Bidokhti, A.A., Ghader, S. and Azadi, M., 2019.** Numerical simulation of oceanic and marine meteorological characteristics of Persian Gulf and Oman Sea using coupled model WRF and ROMS. *Nivar*, 43(106-107), 50-53. DOI: 10.30467/nivar.2019.91474
- Liebich, V., Stehouwer, P. P. and Veldhuis, M., 2012.** Re-growth of potential invasive phytoplankton following UV-based ballast water treatment. *Aquatic Invasions*, 7(1), 29-36. DOI: 10.3391/ai.2012.7.1.004
- Lim, P.T., Leaw, C.P. and Usup, G., 2004.** First incidence of paralytic shellfish poisoning on the east coast of Peninsular Malaysia. *Marine*

- science into the new millennium: new perspectives and challenges. University of Malaya Maritime Research Centre, Kuala Lumpur, pp. 661-667. DOI: 10.1111/pre.12205
- Lim, P.T., Gires, U.S.U.P. and Pin, L.C., 2012.** Harmful algal blooms in Malaysian waters. *Sains Malaysiana*, 41(12), 1509-1515.
- Matsuoka, K., Takano, Y., Kamrani, E., Rezai, H., Sajeevan, T.P. and Al Gheilani, H.M., 2010.** Study on *Cochlodinium polykrikoides* Margalef (Gymnodiniales Dinophyceae) occurring in the Oman Sea and the Persian Gulf from August of 2008 to August of 2009. *Current Development in Oceanography*, 1(3), 153-171.
- MEPC, 2008.** Guidelines for Ballast Water Sampling (G2). International Maritime Organization, London.
- Mirza, R., Mohammadi, M., Faghiri, I., Abedi, E., Fakhri, A., Azimi, A. and Zahed, M.A., 2014.** Source identification of polycyclic aromatic hydrocarbons (PAHs) in sediment samples from the northern part of the Persian Gulf, Iran. *Environmental Monitoring and Assessment*, 186(11), 7387-7398. DOI: 10.1007/s10661-014-3935-y
- Mitbavkar, S. and Anil, A.C., 2006.** Diatoms of the microphytobenthic community in a tropical intertidal sand flat influenced by monsoons: spatial and temporal variations. *Marine Biology*, 148(4), 693-709. DOI: 10.1007/s00227-005-0112-4
- Pereira, N.N., Botter, R.C., Folena, R.D., Pereira, J.P.F.N. and da Cunha, A.C., 2014.** Ballast water: A threat to the Amazon Basin. *Marine Pollution Bulletin*, 84(1-2), 330-338. DOI: 10.1016/j.marpolbul.2014.03.053
- Rey, A., Basurko, O.C. and Rodríguez-Ezpeleta, N., 2018.** The challenges and promises of genetic approaches for ballast water management. *Journal of Sea Research*, 133, 134-145. DOI: 10.1016/j.seares.2017.06.001
- Roziawati, M.R., Leaw, C.P., Lim, H.C., Lee, N., Ishak, I., and Lim, P.T., 2015.** Harmful microalgae assemblage in the aquaculture area of aman island, northern Strait of Malacca. *Malaysian Journal of Science*, 34(1), 20-32. DOI: 10.22452/mjs.vol34no1.3
- Ruiz, G.M., Carlton, J.T., Grosholz, E.D. and Hines, A.H., 1997.** Global invasions of marine and estuarine habitats by non-indigenous species: mechanisms, extent, and consequences. *American zoologist*, 37(6), 621-632. DOI: 10.1093/icb/37.6.621
- Sangpour, M., Ouf, F., Shapouri, M. and Rahimi Bashr, M.R., 2010.** Phytoplankton species in ballast water of two oil tankers in Khark Island. *Journal of Marine Biology*, 1(4), 109-120.
- Saraji, F., 2018.** Establishing a database for phytoplankton species

- from Persian Gulf and Oman Sea, Phase 1: the relevant conducted studies- Iranian Fisheries Science Research Institute. Persian Gulf and Oman Sea Ecology Research Center, Register: NO. 53910.
- Satir, T., 2014.** Ballast water treatment systems: design, regulations, and selection under the choice varying priorities. *Environmental Science and Pollution Research*, 21(18), 10686-10695. DOI:10.1007/s11356-014-3087-1
- Shapoori, M. and Gholami, M., 2014.** Effect of a ballast water treatment system on survivorship of natural populations of marine plankton in Persian Gulf, Iran. *Marketing Science*, 4, 44-48. DOI:10.5923/j.ms.20140402.03
- Shen, L., Xu, H., Guo, X. and Wu, P., 2012.** Oceanography of Skeletonemacostatum harmful algal blooms in the East China Sea using MODIS and QuickSCAT satellite data. *Journal of Applied Remote Sensing*, 6(1): 063529. DOI: 10.1117/1.JRS.6.063529
- Shirowzhan, M., Shanaki, M., Sebt, M.H. and Toosi, H.N., 2016.** Evaluating delay factors in the construction and operation of port operational areas (case study: Shahid Rajaei port complex). *Journal of Fundamental and Applied Sciences*, 8(2), 732-763. DOI:10.4314/jfas.8vi2s.33
- Siang, H., Hussain, R.A., Bhubalan, K. and Orosco, C., 2018.** Ballast Water From Ships Berthed At Major Ports Of Malaysia. *Journal of Sustainability Science and Management*, 13(5), 85-99.
- Sidabutar, T., Srimariana, E.S. and Wouthuyzen, S., 2021.** Phytoplankton species potentially Harmful Algal Blooms (HABs) in Jakarta Bay. In *IOP Conference Series: Earth and Environmental Science*, 744(1), 012077. IOP Publishing. DOI: 10.1088/1755-1315/744/1/012077
- Soler-Figueroa, B.M., Fontaine, D.N., Carney, K.J., Ruiz, G.M. and Tamburri, M.N., 2020.** Characteristics of global port phytoplankton and implications for current ballast water regulations. *Marine Pollution Bulletin*, 155, 111165. DOI: 10.1016/j.marpolbul.2020.111165
- Steichen, J.L., Windham, R., Brinkmeyer, R. and Quigg, A., 2012.** Ecosystem under pressure: Ballast water discharge into Galveston Bay, Texas (USA) from 2005 to 2010. *Marine Pollution Bulletin*, 64(4), 779-789. DOI: 10.1016/j.marpolbul.2012.01.028
- Steichen, J.L., Schulze, A., Brinkmeyer, R., Quigg, A., 2014.** All aboard! A biological survey of ballast water onboard vessels spanning the North Atlantic Ocean. *Marine Pollution Bulletin*, 87(1-2), 201-210. DOI: 10.1016/j.marpolbul.2014.07.058
- Tan, T.H., Leaw, C.P., Leong, S.C.Y., Lim, L.P., Chew, S.M., Teng, S.T. and Lim, P.T., 2016.** Marine micro-

- phytoplankton of Singapore, with a review of harmful microalgae in the region. *Raffles Bulletin of Zoology*, 34, 78-96.
- Teen, L.P., GireS, U.S.U.P. and Pin, L.C., 2012.** Harmful algal blooms in Malaysian waters. *Sains Malaysiana*, 41(12), 1509-1515.
- Tjahjono, A., Bambang, A.N. and Anggoro, S., 2017.** Analysis of heavy metal content of Cd and Zn in ballast water tank of commercial vessels in port of TanjungEmas Semarang, Central Java Province. In *IOP Conference Series: Earth and Environmental Science*, 55(1), 012024. IOP Publishing. DOI: 10.1088/1755-1315/55/1/012024
- Tolian, R., Javadzadeh, N., Sanati, A.M., Mohammadi Roozbahani, M. and Noorinejad, M., 2020.** Investigating the Plankton Diversity and Density in ballast tanks of ships entering the Persian Gulf (Bushehr Port). *Journal of Marine Biology*, 12(1), 63-82.
- Tomas, C.R., 1997.** Identifying marine phytoplankton. Elsevier.
- Tsolaki, E. and Diamadopoulos, E., 2010.** Technologies for ballast water treatment: a review. *Journal of Chemical Technology and Biotechnology*, 85(1), 19-32. DOI: 10.1002/jctb.2276
- Van Slooten, C., 2021.** Ballast water treatment system testing: assessing novel treatments and validating compliance methods. University of Groningen, 165 P. DOI: 10.33612/diss.172082815
- Vieira, D.P., de Mello, P.C., Dotta, R. and Nishimoto, K., 2018.** Experimental investigation on the influence of the liquid inside the tanks in the wave behavior of FLNG vessels in side-by-side offloading operations. *Applied Ocean Research*, 74, 28-39. DOI: 10.1016/j.apor.2018.02.019
- Wang, Z. and Liu, Y., 2021.** Research on the Impact of Policy Incentives on the Implementation of Ship ECA Policies. In *IOP Conference Series: Earth and Environmental Science* (Vol. 638, No. 1, p. 012004). IOP Publishing. DOI:10.1088/1755-1315/638/1/012004
- Werschkun, B., Banerji, S., Basurko, O.C., David, M., Fuhr, F., Gollasch, S. and Höfer, T., 2014.** Emerging risks from ballast water treatment: The run-up to the International Ballast Water Management Convention. *Chemosphere*, 112, 256-266. DOI: 10.1016/j.chemosphere.2014.03.135
- Wu, H., Chen, C., Wang, Q., Lin, J. and Xue, J., 2017.** The biological content of ballast water in China: A review. *Aquaculture and Fisheries*, 2(6), 241-246. DOI: 10.1016/j.aaf.2017.03.002.